

Estimated Vaccination Coverage by 24 Months of Age, by Selected Vaccines and Doses for the United States, Department of Health and Human Services (HHS) Region I, and Vermont– National Immunization Survey, 2014

Vaccine and doses	United States		HHS Region I*		Vermont	
	%	(95% CI)	%	(95% CI)	%	(95% CI)
≥ 4 doses DTaP	80.6	(±1.3)	86.5	(±3.1)	85.8	(±5.4)
≥ 3 doses Poliovirus	92.5	(±0.9)	93.1	(±2.4)	91.0	(±4.2)
≥ 1 dose MMR	89.5	(±1.0)	93.2	(±2.2)	90.6	(±4.0)
≥ 3 doses Hib	91.9	(±0.9)	95.2	(±2.0)	94.6	(±3.6)
≥ 3 doses HepB	90.9	(±0.9)	92.4	(±2.2)	88.8	(±4.5)
≥ 1 dose Varicella	88.9	(±1.0)	92.0	(±2.5)	83.7	(±5.6)
≥ 4 doses PCV	80.9	(±1.3)	85.8	(±3.2)	84.3	(±5.5)
≥ 1 dose HepA	81.6	(±1.2)	83.7	(±3.2)	72.9	(±6.6)
Rotavirus†	71.7	(±1.6)	79.6	(±3.7)	76.6	(±6.3)
≥ 2 doses Influenza	52.8	(±1.6)	71.2	(±4.0)	63.3	(±7.2)
Combined Series§	36.2	(±1.6)	47.2	(±4.5)	40.7	(±7.4)

Abbreviations: CI = confidence interval; DTaP = diphtheria, tetanus toxoids, and acellular pertussis vaccine (includes children who might have been vaccinated with diphtheria and tetanus toxoids vaccine, or diphtheria, tetanus toxoids, and pertussis vaccine); MMR = measles, mumps, and rubella vaccine; Hib = *Haemophilus influenzae* type b vaccine; HepB = hepatitis B vaccine; PCV = pneumococcal conjugate vaccine; HepA = hepatitis A vaccine.

* HHS Region I includes Connecticut, Maine, Massachusetts, New Hampshire, Rhode Island, and Vermont.

† Rotavirus vaccine includes ≥2 or ≥3 doses, depending on the product type received (≥2 doses for Rotarix [RV1] or ≥3 doses for RotaTeq [RV5]).

§ Combined series includes ≥ 4 doses of DTaP, ≥ 3 doses of poliovirus vaccine, ≥ 1 dose of MMR, ≥ 3 doses of Hib, ≥ 3 doses of HepB, ≥ 1 dose of varicella vaccine, ≥ 4 doses of PCV, ≥ 1 dose of HepA, ≥ 2 or ≥ 3 doses of rotavirus vaccine (depending on product type), and ≥ 2 doses of influenza vaccine.