## 2011 Vermont Youth Risk Behavior Survey Report

## **Report for Participating Schools in Addison County**

Every other year, since 1993, the Department of Health Division of Alcohol and Drug Abuse Programs and the Department of Education Student Health and Learning team sponsor the Vermont Youth Risk Behavior Survey (YRBS). The YRBS measures the prevalence of behaviors that contribute to the leading causes of death, disease, and injury among youth.

Participation by schools and individual students is voluntary. To protect student privacy, the questionnaire and survey process are anonymous. It is impossible to identify an individual student's responses.

### **Survey Changes in 2011**

Before 2011, students in grades 8-12 took the Vermont Youth Risk Behavior Survey. In 2011, we conducted two surveys: a **high school survey** of students in grades 9-12, and a **middle school survey** of students in grades 6-8.

**All results in the high school survey report are for grades 9-12 only.** For the questions on the 2011 high school survey, we recalculated results from previous years for grades 9-12 for comparative purposes. Those results are at the end of this report.

### **About this Report**

The results appear in data tables by gender and grade unless otherwise noted. All results are percentages of students who responded affirmatively. Percentages were suppressed with an asterisk (\*) if five or fewer students reported affirmatively, either overall or by grade or gender.

### **Thank You**

We thank the principals and superintendents who chose to participate, the teachers and school staff who administered or in other ways supported the survey, and the students who took the time and effort to share a piece of their lives with us. We also thank the Centers for Disease Control and Prevention, Division of Adolescent and School Health for sponsoring the survey through a cooperative agreement with the Vermont Department of Education, and the CDC and Westat for technical assistance.

## **Table of Contents**

BACKGROUND INFORMATION	
How to Use the YRBS	
A Word of Caution	2
About the Surveys	5
HIGH SCHOOL SURVEY RESULTS	15
Sample Demographics	16
Personal Safety	
Alcohol, Tobacco, and Other Drug Use	
Attitudes and Perceptions about Alcohol, Cigarette, and Marijuana Use	
Sexual Behavior and Orientation	
Body Image	
Nutrition	
Physical Activity	
Youth Assets	
7041710000	
MIDDLE SCHOOL SURVEY RESULTS	39
Sample Demographics	
Personal Safety	
Alcohol, Tobacco, and Other Drug Use	
Attitudes and Perceptions about Alcohol, Cigarette, and Marijuana Use	
Body Image	
Nutrition	
Physical Activity	
Youth Assets	
RECALCULATED HIGH SCHOOL RESULTS	57

### How to Use the YRBS

The YRBS can detect changes in risk behaviors over time and identify differences among ages, grades, and genders. With these data, we can focus prevention efforts and determine whether school policies and community programs are having the intended effect on student behaviors.

Think of the YRBS as a tool for starting discussions, for educating the community, for planning and evaluating programs, and for comparing Vermont students with other students nationwide.

- Start the Conversation: Use the YRBS to begin a conversation with teens about the personal choices they make or about the health of their community. Ask them if the results accurately reflect what they see happening around them. How do they explain the results? From their perspective, what is or is not working? How would they promote healthy behaviors?
- Increase Awareness: The YRBS provides an opportunity to break through "denial" and make community members aware of the risks that their young people face. It can also dispel myths and correct misinformation about the "average teenager." The YRBS can accentuate the positive and celebrate the fact that many students are abstaining from behaviors that endanger their health and their ability to succeed.
- Plan and Evaluate Programs: The YRBS can serve as the basis of a community needs assessment. It can help identify strengths and weaknesses in communities, and can inform strategies to address those weaknesses.

**Remember to Look at the Positive Side:** In most cases, the majority of adolescents are NOT engaging in risky behaviors. Although most of the charts examine the prevalence of risk behaviors, please do not forget about the percentage of adolescents who are NOT engaging in these behaviors.

### A Word of Caution

The YRBS represents the most complete and most recent information available about risk behaviors among Vermont students. However, the YRBS has some limitations that you should keep in mind when interpreting the results.

- Data Quality: We take numerous precautions to ensure the reliability and validity of
  the results. The Centers for Disease Control and Prevention (CDC) carefully
  designed and thoroughly tested the questionnaire. The anonymous survey
  encourages students to be honest and forthright. We run over 100 consistency
  checks on the data to exclude careless, invalid, or logically inconsistent answers.
  These precautions can reduce some sources of error, but not all.
- Comparing Supervisory Unions/School Districts to Each Other and to the State: It is natural to want to know how individual supervisory unions or school districts compare to the state overall or to each other. We urge caution in making such comparisons because the statewide results are weighted and the local results are not. As a result, it is possible that apparent differences, especially small differences, are due to demographic characteristics rather than to true differences in prevalence. Furthermore, small differences may not be statistically different and may simply be a function of normal sampling error.
- What, not Why: The YRBS can indicate what students are doing. It can also suggest which groups of students are more likely to engage in these behaviors. However, the survey does not answer the most important question: Why are they doing it?

Feeling safe – whether at school, in a car, or in a relationship – fosters positive adolescent development. The high school and middle school surveys included questions on fighting, bullying, seat belt use, safety at school, helmet use, and suicidality. The high school survey included additional questions on driving behaviors, dating violence, and self harm.

- Physical fighting is a marker for problem behaviors<sup>1</sup> and is associated with serious injury.<sup>2,3</sup> Abuse by an intimate partner and forced sex are associated with negative psychosocial outcomes, poor mental health outcomes, and other risk behaviors among both males and females.<sup>4,5,6</sup>
- Both being a bully and being victimized by bullies have been increasingly recognized as health problems for children because of their association with a range of problems, including poor psychological adjustment,<sup>7,8</sup> poor academic achievement,<sup>8</sup> and violent behavior.<sup>9</sup>
- Motor vehicle crash injuries are the leading cause of death among youth ages 15 to 19 in the U.S.<sup>10</sup> Proper use of safety belts reduces the risk of fatal injury to front seat passengers by 45% and risk of moderate to critical injury by 50%.<sup>11</sup>
- Bicycle helmets are 85% to 88% effective at reducing the impact of head and brain injuries due to bicycle crashes. 12 Despite this, less than one quarter of bicyclists wear helmets. 12
- In 2009, alcohol use was associated with nearly four in ten motor vehicle-related fatalities nationwide and in Vermont.<sup>13</sup> Additionally, research examining drugs other than alcohol indicates that marijuana is the most prevalent drug detected in impaired drivers, fatally injured drivers, and motor vehicle crash victims nationwide.<sup>14</sup>

\_\_\_\_\_

## Alcohol, Tobacco, and Other Drugs

Early use of alcohol, tobacco, and other drugs is associated with myriad problems later in life. The high school and middle school surveys asked about tobacco, alcohol, marijuana, inhalant, and prescription drug use. The surveys also asked the age at which students first used alcohol, cigarettes, and marijuana. The high school survey asked additional questions on other drug use, such as heroin, methamphetamines, and hallucinogens. Both surveys asked about ease of availability of alcohol, cigarettes, and marijuana, peer disapproval of use of these substances, perceived parental disapproval of use, and the risk of harm associated with use of these substances.

- Alcohol use is a major contributing factor in about half of all homicides and sexual assaults,<sup>15</sup> and about one-third of all motor vehicle crash fatalities.<sup>16</sup> Heavy drinking among youth has been linked to violence, academic and job problems, suicidal behavior, trouble with law enforcement authorities, risky sexual behavior, and use of cigarettes, marijuana, cocaine, and other illegal drugs.<sup>17,18</sup>
- Tobacco use is the single most preventable cause of death in the United States,<sup>19</sup> contributing to more than one of every five deaths.<sup>20</sup> Cigarette smoking increases the risk of heart disease, chronic obstructive pulmonary disease, acute respiratory illness, stroke, and cancers of the lung, larynx, oral cavity, pharynx, pancreas, and cervix.<sup>19</sup>
- Marijuana use is associated with smoking-related respiratory damage, temporary short-term memory loss, decreased motivation, and psychological dependence.<sup>21</sup> Other reactions include feelings of distrust, anxiety, or depression.<sup>21</sup> In Vermont, more teens enter treatment with a primary diagnosis for marijuana dependence than all other illicit drugs combined.<sup>22</sup>
- Other drug use is related to suicide, early unwanted pregnancy, school failure, delinquency, and transmission of sexually transmitted diseases (STDs), including HIV infection.<sup>23</sup>
- Multi-year results from the Monitoring the Future survey indicate that the prevalence of marijuana use among youth declines as the percentage of youth expressing disapproval of marijuana increases.<sup>24</sup> As perception of harm of using alcohol and other drugs decreases, there is a tendency for use to increase.<sup>24</sup> Increased use is also associated with the perception that substances are readily available, regardless of whether the perception is accurate.<sup>24</sup>

### **Sexual Behavior and Orientation**

Experimentation with sexual behaviors and orientation often starts during adolescence. The high school survey asked about age of first sexual intercourse, frequency, sexual partners, sexual orientation, alcohol and drug use related to sexual intercourse, contraceptive use, and HIV testing. The middle school survey did not ask any questions about sex.

- Early sexual activity and having multiple sexual partners are associated with an increased risk of unwanted pregnancy, sexually transmitted diseases (STDs) including HIV infection,<sup>25</sup> and negative effects on social and psychological development.<sup>26</sup> Alcohol and drug use may serve as predisposing factors for initiation of sexual activity.<sup>27</sup>
- Of the nearly 19 million new cases of STDs per year in the United States, almost half are among youth ages 15-24.<sup>28</sup> STDs may result in infertility and facilitation of HIV transmission and may have an adverse effect on pregnancy outcomes and maternal and child health.<sup>26</sup> Besides abstinence, condom use is currently the most effective means of preventing sexual transmission of HIV and other STDs.<sup>29</sup>
- Although many lesbian, gay, bisexual, and transgender adolescents lead happy and healthy lives, others face tremendous challenges to growing up physically and mentally healthy.
   Compared to heterosexual youth, lesbian, gay, bisexual, and transgender youth are at higher risk for depression, tobacco, alcohol and other drug use, suicide, and unhealthy sexual behaviors.<sup>30</sup>

## **Body Image**

Negative feelings about weight and body image often develop in adolescence. The high school and middle school surveys included questions about weight perception and weight control. The high school survey asked for students' height and weight to calculate body mass index, and about use of artificial tanning devices.

- There are more than three times as many overweight children and adolescents in the U.S. than there were in 1980.<sup>31</sup> Obesity in childhood and adolescence is associated with negative psychological and social consequences and adverse health outcomes, including type 2 diabetes, obstructive sleep apnea, hypertension, dyslipidemia, and metabolic syndrome.<sup>32</sup> Overweight and obesity acquired during childhood or adolescence may persist into adulthood.<sup>33</sup> Approximately 400,000 deaths a year in the United States are currently associated with overweight and obesity.<sup>20</sup>
- However, overemphasis on slenderness during adolescence may contribute to eating disorders such as anorexia nervosa and bulimia.<sup>34,35</sup> Studies have shown high rates of body dissatisfaction and dieting among adolescent females, with many engaging in unhealthy weight control behaviors such as fasting and self-induced vomiting that can lead to abnormal physical and psychological development.<sup>36,37</sup>
- Indoor tanning increases risk for melanoma and non-melanoma cancers. In the U.S., the
  incidence of melanoma is increasing more rapidly than that of any other cancer, particularly
  among girls and women between the ages of 15 and 39. Some evidence suggests that
  repeated UV irradiation, and the use of indoor tanning beds specifically, may have important
  behavioral consequences, including mood changes, pain, and physical dependency.<sup>38</sup>

## **Nutrition and Physical Activity**

Nutritious eating and physical activity are two cornerstones of healthy adolescent development. The high school and middle school surveys asked about breakfast consumption, physical activity, physical education classes, and television, computer, and video game use. The high school survey also asked about consumption of fruits, vegetables, soda, and sugar-sweetened beverages.

- Fruits and vegetables are good sources of complex carbohydrates, vitamins, minerals, and
  other substances that are important for good health. Dietary patterns with higher intakes of
  fruits and vegetables are associated with a variety of health benefits, including a decreased
  risk for some types of cancer.<sup>39-43</sup>
- In recent years, soft drink consumption has significantly increased among children and adolescents. Consumption of sugar-sweetened drinks, including soft drinks, appears to be associated with an increased risk for being overweight in children.
- Regular physical activity builds and maintains healthy bones and muscles, controls weight, builds lean muscle, reduces fat, reduces feelings of depression and anxiety. It also decreases the risk of dying prematurely, dying of heart disease, and developing diabetes, colon cancer, and high blood pressure. The U.S. Department of Health and Human Services recommends that young people ages 6–17 participate in at least 60 minutes of physical activity every day.
- By 12th grade, more than half of female students in the U.S. do not participate in vigorous physical activity regularly. School physical education classes can increase adolescent participation in physical activity and help adolescents develop the knowledge, attitudes, and skills they need to engage in lifelong physical activity.<sup>47-50</sup>
- Television viewing is the principal sedentary leisure time behavior in the U.S. Studies have shown that television viewing in young people is related to obesity<sup>51</sup> and violent or aggressive behavior.<sup>52-54</sup> Using the computer for fun and playing video games have become increasingly common sedentary leisure time activities among young people as well.

#### **Measures of Youth Assets**

Adolescent achievement requires sources of positive influence. The high school and middle school surveys asked about school and community connectivity as well as parental conversations about school. The high school survey asked additional questions about volunteerism and grades earned in school.

- Above-average school performance is one of many developmental assets, or factors promoting positive development. Studies have shown that students who get higher grades in school are less likely to use cigarettes, alcohol, or marijuana, and are more likely to postpone sexual intercourse.<sup>55</sup>
- One of the strongest predictors of students' success in school is the extent to which their parents stay involved with their schoolwork.<sup>56</sup> A national study of adolescent health found that youth who reported a "connectedness" to their parents/family and school were the least likely to engage in risky behaviors.<sup>57</sup> Parental expectations regarding school achievement were also associated with lower levels of risk behaviors.<sup>57</sup>
- Research shows that involvement in constructive, supervised extracurricular
  activities is associated with reduced likelihood of involvement in risky behaviors such
  as school failure, drug use, and delinquency.<sup>58</sup> In addition, evidence is emerging that
  students who participate in such activities are also more likely to engage in other
  "thriving" behaviors.<sup>59</sup>
- Youth are not simply objects of adult efforts to modify their behaviors. Rather, if given the opportunities, they can make significant contributions to their families, schools, and communities. Adolescents, especially, need to exercise decision-making power in as many settings as is practical, so that they can develop into competent adults. Schools are a natural setting for youth to share in decisions that affect their lives.

### References

1. Sosin, D.M., Koepsell, T.D., Rivara, F.P., Mercy, J.A. Fighting as a marker for multiple problem behaviors in adolescents. <u>Journal of Adolescent Health</u> 16(3):209-215, 1995.

- 2. Borowsky, I.W., Ireland, M. Predictors of future fight-related injury among adolescents. <u>Pediatrics</u> 113(3 pt 1):530-536, 2005.
- 3. Pickett, W., Craig, W., Harel, Y., et al. Cross-national study of fighting and weapon carrying as determinants of adolescent injury. Pediatrics 116(6):e855-863, 2005.
- 4. Roberts, T.A., Klein, J.D., Fisher, S. Longitudinal effect of intimate partner abuse and high-risk behavior among adolescents. <u>Archives of Pediatrics and Adolescent Medicine</u> 157(9):875-881, 2003.
- 5. Ackard, D.M., Neumark-Sztainer, D. Date violence and date rape among adolescents: association with disordered eating behaviors and psychological health. <u>Child Abuse and Neglect</u> 26(5):455-473, 2002.
- 6. Howard, D.E., Wang, M.Q. Psychosocial correlates of U.S. adolescents who report a history of forced sexual intercourse. <u>Journal of Adolescent Health</u> 36(5):372-379, 2005.
- 7. Juvonen, J., Graham, S., Schuster, M.A. Bullying among young adolescents: the strong, the weak, and the troubled. Pediatrics 112(6 pt 1): 1231-1237, 2003.
- 8. Spivak, H., Prothrow-Stith, D. The need to address bullying-an important component of violence prevention. <u>JAMA</u> 285(16):2131-2132, 2001.
- 9. Nansel, T.R., Overpeck, M., Pilla, R.S., et al. Bullying behaviors among U.S. youth: prevalence and association with psychological adjustment. <u>JAMA</u> 285(16):2094-2100, 2001.
- 10. Centers for Disease Control and Prevention. Web-based Injury Statistics Query and Reporting System (WISQARS). National Center for Injury Prevention and Control, Centers for Disease Control and Prevention. Online: www.cdc.gov/injury/wisqars/index.html
- National Highway Traffic Safety Administration. Traffic safety facts: occupant protection, 2007.
   Online: www-nrd.nhtsa.dot.gov/Pubs/810991.PDF
- 12. National Highway Traffic Safety Administration. Traffic safety facts: bicycle helmet use laws, 2008. Online: www.nhtsa.dot.gov/people/injury/TSFLaws/PDFs/810886.pdf
- National Highway Traffic Safety Administration. Traffic safety facts: state alcohol-impaired driving estimates, 2009. Online: www-nrd.nhtsa.dot.gov/Pubs/811398.pdf
- Jones, R.K., Shinar, D., Walsh, J.M. <u>State of Knowledge of Drug-Impaired Driving</u>. National Highway Traffic Safety Administration Technical Report DOT HS 809 642. Washington, DC: U.S. Department of Transportation, 2003.
- 15. Abbey, A., Zawacki, T., Buck, P.O., et al. Alcohol and sexual assault. <u>Alcohol Research and Health</u> 25(1):43-51, 2001.
- 16. Miller, J.W., Naimi, T.S., Brewer, R.D., Jones, S.E. Binge drinking and associated health risk behaviors among high school students. <u>Pediatrics</u> 119(1):76-85, 2007.

### References

17. National Research Council and Institute of Medicine (2004). Reducing Underage Drinking: A Collective Responsibility. Committee on Developing a Strategy to Reduce and Prevent Underage Drinking, Richard J. Bonnie and Mary Ellen O'Connell, Editors. Board on Children, Youth, and Families, Division of Behavioral and Social Sciences and Education. Washington, DC: The National Academies Press.

- 18. U.S. Department of Health and Human Services. <u>The Surgeon General's Call to Action to Prevent and Reduce Underage Drinking</u>. U.S. Department of Health and Human Services, Office of the Surgeon General, 2007.
- U.S. Department of Health and Human Services. <u>The Health Consequences of Smoking: A Report of the Surgeon General</u>.
   U.S. Department of Health and Human Services; Centers for Disease Control and Prevention; National Center for Chronic Disease Prevention and Health Promotion: Office on Smoking and Health, 2004.
- 20. Mokdad, A.H., Marks, J.S., Stroup, D.F., Gerberding, J.L. Actual causes of death in the United States, 2000. JAMA 291(10):1238-1245, 2004.
- 21. National Institute on Drug Abuse. <u>Research Report Series: Marijuana Abuse</u> (NIH Publication 05-3859). Bethesda, MD: National Institute on Drug Abuse, 2002.
- 22. Vermont Substance Abuse Treatment Information System. Data online at: www.healthvermont.gov/adap/clearinghouse/documents/AdolescentsbySAandFY.pdf
- 23. Newcomb, M.D., Locke T. Health, social, and psychological consequences of drug use and abuse. In: Epidemiology of Drug Abuse (Z. Sloboda, ed.). Springer U.S., 2006.
- 24. Johnston, L., O'Malley, P., Bachman, J. G., Shulenberg, J.E. <u>National Survey Results on Drug Use From the Monitoring the Future Study, 1975-2007, Volume I: Secondary School Students</u> (NIH Publication No. 08-6418A). Bethesda, MD: National Institute of Drug Abuse, 2008.
- 25. Abma JC, Martinez GM, Copen CE. Teenagers in the United States: Sexual activity, contraceptive use, and childbearing, National Survey of Family Growth 2006-2008. National Center for Health Statistics. Vital Health Stat 23 (30). 2010.
- 26. Centers for Disease Control and Prevention. Fact sheet on STDs and pregnancy. Online: www.cdc.gov/std/pregnancy/STDs-and-pregnancy-fact-sheet.pdf
- 27. Cavazos-Rehg, P.A., Krauss, M.J., Spitznagel, E.L., et al. Substance use and the risk for sexual intercourse with and without a history of teenage pregnancy among adolescent females. <u>Journal of Studies on Alcohol and Drugs</u> 72(2): 194-198, 2011.
- 28. Gavin, L., MacKay, A.P., Brown, K., et al. Sexual and reproductive health of persons aged 10-24 years United States, 2002-2007. MMWR Surveillance Summaries 58(6): 1-58, 2009.
- 29. Joint United Nations Programme on HIV/AIDS (UNAIDS). Fast Facts about HIV Prevention. Online at: www.unaids.org/en/media/unaids/contentassets/dataimport/pub/basedocument/2008/20080501\_fastf acts prevention en.pdf
- 30. Kann, L., Olsen, E.O., McManus, T., et al. Sexual Identity, Sex of Sexual Contacts, and Health-Risk Behaviors Among Students in Grades 9-12 Youth Risk Behavior Surveillance, Selected Sites, United States, 2001-2009. MMWR Early Release 60(7): 1-133, 2011.
- 31. Ogden, C.L., Carrol, M.D. Prevalence of obesity among children and adolescents: United States, trends 1963-1965 through 2007-2008. National Center for Health Statistics Health E-Stats, June 2010.

### References

32. Daniels, S.R., Arnett, D.K., Eckel, R.H., et al. Overweight in children and adolescents: pathophysiology, consequences, prevention, and treatment. Circulation 111(15):1999-2012, 2005.

- 33. Wright, C.M., Parker, L., Lamont, D., Craft, A.W. Implications of childhood obesity for adult health: findings from thousand families cohort study. <u>British Medical Journal</u> 323(7324):1280-1284, 2001.
- 34. Tremblay, L., Lariviere, M. The influence of puberty onset, body mass index, and pressure to be thin on disordered eating behaviors in children and adolescents. <u>Eating Behaviors</u> 10(2):75-83, 2009.
- 35. Mitchell, J.E., Eckert, E.D. Scope and significance of eating disorders. <u>Journal of Consulting Clinical Psychology</u> 55:628-634, 1987.
- 36. Neumark-Sztainer, D., Hannan, P.J. Weight-related behaviors among adolescent girls and boys: results from a national survey. <u>Archives of Pediatric and Adolescent Medicine</u> 154(6):569-577, 2000.
- 37. Neumark-Sztainer, D., Story, M., Hannan, P.J., et al. Weight-related concerns and behaviors among overweight and nonoverweight adolescents: implications for preventing weight-related disorders.

  <u>Archives of Pediatric and Adolescent Medicine</u> 156(2):171-178, 2002.
- 38. Fisher, D.E. and James, W.D. Indoor tanning science, behavior, and policy. <u>New England Journal of Medicine</u> 363:901-903, 2010.
- 39. Key, T.J., Schatzkin, A., Willet, W.C., et al. Diet, nutrition, and the prevention of cancer. <u>Public Health Nutrition</u> 7(1A):187-200, 2004.
- National Cancer Institute. <u>5 A Day for Better Health Program</u> (NIH Publication 01-5019). Betheseda, MD, 2001.
- 41. Kavey, R.E., Daniels, S.R., Lauer, R.M., et al. American Heart Association guidelines for primary prevention of atherosclerotic cardiovascular disease beginning in childhood. <u>Journal of Pediatrics</u> 142(4):368-372, 2003.
- 42. Terry, P., Terry, J.B., Wolk, A. Fruit and vegetable consumption in the prevention of cancer: an update. <u>Journal of Internal Medicine</u> 250(4):280-290, 2001.
- 43. Van Duyn, M.A., Pivonka, E. Overview of the health benefits of fruit and vegetable consumption for the dietetics professional: selected literature. <u>Journal of the American Dieticians Association</u> 100(12):1511-1521, 2000.
- 44. Malik, V.S., Schulze, M.B., Hu, F.B. Intake of sugar-sweetened beverages and weight gain: a systematic review. American Journal of Clinical Nutrition 84(2):274-288, 2006.
- 45. Ludwig, D.S., Peterson, K.E., Gortmaker, S.L. Relation between consumption of sugar-sweetened drinks and childhood obesity: a prospective, observational analysis. <u>Lancet</u> 357(9255):505-508, 2001.
- 46. Physical Activity Guidelines Advisory Committee. <u>Physical Activity Guidelines Advisory Committee</u> <u>Report, 2008</u>. Washington, D.C.: U.S. Department of Health and Human Services, 2008.
- 47. Durant, N., Harris, S.K., Doyle, S., et al. Relation of school environment and policy to adolescent physical activity. Journal of School Health 79(4):153-159, 2009.
- 48. McKenzie, K.L., Li, D., Derby, C.A., et al. Maintenance of effects of the CATCH Physical Education Program: results from the CATCH-ON Study. <u>Health Education & Behavior</u> 30(4):447-462, 2003.

### References

49. U.S. Department of Health and Human Services and U.S. Department of Education. Promoting better health for young people through physical activity and sports. 2000. Online: www.cdc.gov/HealthyYouth/physicalactivity/promoting health/pdfs/ppar.pdf

- 50. Center for Disease Control and Prevention. Guidelines for school and community programs to promote lifelong physical activity among young people. Morbidity and Mortality Weekly Report 46 (No. RR-6):1-36, 1997.
- 51. Zabinski, M.F., Norman, G.J., Sallis, J.F., et al. Patterns of sedentary behavior among adolescents. Health Psychology 26(1):113-120, 2007.
- 52. Crespo, C.J., Smit, E., Troiano, R.P., et al. Television watching, energy intake, and obesity in U.S. children: results from the third National Health and Nutrition Examination Survey, 1988-1994. <u>Archives of Pediatric and Adolescent Medicine</u> 155(3):360-365, 2001.
- 53. Kaur, H., Choi, W.S., Mayo, M.S., Harris, K.J. Duration of television watching is associated with increased body mass index. Journal of Pediatrics 143(4):506-511, 2003.
- 54. Kuntsche, E., Pickett, W., Overpeck, M., et al. Television viewing and forms of bullying among adolescents from eight countries. Journal of Adolescent Health 39(6):908-915, 2006.
- 55. Resnick, M.D., Bearman, P.S., Blum, R.W., et al. Protecting adolescents from harm. Findings from the National Longitudinal Study on Adolescent Health. <u>JAMA</u> 278(10):823-832, 1997.
- 56. Fan, X., Chen, M. Parental involvement and students' academic achievement: a meta-analysis. <u>Educational Psychology Review</u> 13(1):1-22, 2001.
- 57. U.S. Council of Economic Advisors. Teens and their parents in the 21<sup>st</sup> century: An examination of trends in teen behavior and the role of parental involvement. 2000. Online: http://clinton3.nara.gov/WH/EOP/CEA/html/Teens\_Paper\_Final.pdf
- 58. Fredricks, J.A., Eccles, J.S. Is extracurricular participation associated with beneficial outcomes? <u>Developmental Psychology</u> 42(4):698-713, 2006.
- 59. Scales, P.C., Benson, P.L., Leffert, N., Blyth, D.A. Contribution of developmental assets to prediction of thriving among adolescents. Applied Developmental Science 4(1):27-46, 2000

# 2011 Vermont High School Youth Risk Behavior Survey

# **Results for Addison County**

### Participation

	Grade			Sex		
9	10	11	12	Female	Male	All
292	332	380	284	608	652	1271

For the high school survey, students self-reported the town in which they live. We assigned students to counties based on this town code. Students who did not provide a useable town code are not included in the county-level reports.

### **Participating Schools:**

Middlebury Union High School Mount Abraham Union MS/HS Patricia A. Hannaford Career Center Vergennes Union High School

# Demographics

Age

	2011
14 or younger	15%
15	26%
16	27%
17	25%
18 or older	7%

## Race and Ethnicity

	2011
White non-Hispanic	91%
Racial or ethnic minority	9%

### Mother's Education

	2011
High school or less	28%
Some college	13%
College graduate	35%
Graduate or professional degree after college	16%
Not sure	7%

Percent of students who:

	All	Grade			All Grade Sex		<
	2011	9	10	11	12	Female	Male
Ride bicycles	74%	81%	83%	70%	63%	69%	79%

Of students who ride bicycles, percent who wear helmets:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Always	15%	17%	15%	13%	17%	15%	15%
Most of the time	13%	19%	10%	14%	7%	13%	12%
Sometimes	12%	13%	10%	12%	12%	11%	13%
Rarely or never	60%	51%	65%	61%	63%	61%	60%

Percent of students who wear seatbelts when riding in a car:

	All	Grade				Sex		
	2011	9	10	11	12	Female	Male	
Always	63%	63%	61%	65%	63%	68%	58%	
Most of the time	23%	26%	27%	21%	19%	21%	25%	
Sometimes	8%	6%	7%	7%	11%	6%	9%	
Rarely or never	6%	6%	5%	7%	7%	5%	7%	

Percent of students who, in the past 30 days:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Rode in a car or other vehicle driven by someone who had been drinking alcohol	19%	20%	17%	21%	20%	18%	20%
Drove a car or other vehicle after drinking	5%	3%	3%	6%	9%	4%	6%
Rode in a car or other vehicle driven by someone who had been smoking marijuana	21%	11%	17%	25%	31%	20%	22%
Drove a car or other vehicle after smoking marijuana	10%	4%	6%	12%	20%	9%	12%

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Carried a weapon such as a gun, knife or club on school property, past 30 days	12%	11%	10%	14%	14%	6%	19%
Did not go to school because they felt unsafe, past 30 days	3%	4%	3%	4%	3%	3%	4%
Were threatened or injured with a weapon at school, past 12 months	5%	8%	6%	3%	2%	2%	7%
Were in a physical fight, past 12 months	23%	30%	24%	23%	15%	17%	29%
Were in a physical fight on school property, past 12 months	10%	12%	12%	10%	5%	5%	13%
Were hit, slapped, or physically hurt on purpose by a boyfriend or girlfriend, past 12 months	7%	5%	7%	8%	7%	6%	7%
Were physically forced to have sexual intercourse, ever	5%	3%	5%	6%	8%	8%	3%

Percent of students who:

	All		Gra	ade		Sex	
	2011	9	10	11	12	Female	Male
Were bullied, past 30 days	16%	20%	17%	16%	11%	18%	14%
Bullied someone else, past 30 days	17%	19%	14%	17%	16%	15%	18%
Were electronically bullied, past 12 months	13%	14%	12%	16%	9%	18%	8%
Caused purposeful harm to themselves such as by cutting or burning, past 12 months	13%	10%	14%	13%	14%	19%	7%
Felt sad or hopeless almost every day for two weeks or more in a row, past 12 months	17%	13%	19%	19%	18%	23%	12%
Made a suicide plan, past 12 months	7%	8%	9%	5%	7%	9%	6%
Made a suicide attempt, past 12 months	3%	3%	4%	2%	2%	3%	3%

For the purposes of the Vermont YRBS, bullying was described as occurring when, on many occasions, a student or group of students say or do unpleasant things to another student to make fun of, tease, embarrass, or scare him/her, or purposefully exclude him/her. Bullying can occur before, during, or after the school day; on school property; on a school bus; or at a school-sponsored activity. It is not bullying when two students of about the same strength and power argue or fight, or when teasing is done in a friendly way. Electronic bullying includes through e-mail, chat rooms, instant messaging, Web sites, or texting.

	All		Gra	Sex	ĸ		
	2011	9	10	11	12	Female	Male
Ever drank alcohol other than a few sips	60%	43%	51%	67%	78%	59%	60%
Had first drink other than a few sips before age 13	16%	19%	16%	16%	15%	11%	21%
Drank alcohol, past 30 days	32%	20%	27%	36%	47%	32%	32%
Had five or more drinks of alcohol in a row, past 30 days	17%	9%	14%	18%	27%	15%	19%
Drank alcohol on 3 to 9 of the past 30 days	11%	6%	8%	12%	18%	10%	11%
Drank alcohol on 10 or more of the past 30 days	3%	2%	3%	3%	6%	2%	5%
Binged on 3 or more of the past 30 days	7%	2%	7%	7%	11%	4%	9%
Drank alcohol on school property, past 30 days	2%	3%	3%	*	3%	2%	3%

<sup>\*</sup> Suppressed due to small cell sizes (≤ 5 students)

Among students who drank in the past 30 days:

Usual type of beverage consumed, past 30 days

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Liquor, such as rum, gin, vodka, or whiskey	46%	49%	49%	41%	48%	49%	44%
Beer	32%	28%	29%	32%	35%	20%	43%
Other alcoholic beverages, such as Smirnoff Ice, Bacardi Silver, Mike's Hard Lemonade, or Twisted Tea	17%	17%	16%	22%	13%	24%	10%
Wine	5%	*	*	*	*	7%	3%

Usual source of alcohol, past 30 days

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Someone gave it to me	49%	38%	53%	53%	47%	51%	47%
Gave someone money to buy it for me	26%	*	22%	28%	33%	25%	26%
Got it or stole it from home	20%	45%	24%	13%	12%	21%	19%
Purchased from a store, liquor store; or restaurant	5%	*	*	*	*	*	7%
Stole it from a store or restaurant	*	*	*	*	*	*	*

<sup>\*</sup> Suppressed due to small cell sizes (≤ 5 students)

	All		Gra	ade		Sex	
	2011	9	10	11	12	Female	Male
Ever smoked a whole cigarette	25%	15%	20%	28%	38%	24%	26%
Smoked before age 13	8%	6%	8%	10%	8%	6%	10%
Smoked cigarettes, past 30 days	13%	7%	10%	13%	22%	14%	12%
Smoked on 20 or more of the past 30 days	4%	*	3%	4%	10%	5%	4%
Smoked every day, past 30 days	4%	*	2%	3%	9%	4%	3%
Used chewing tobacco, past 30 days	7%	5%	6%	8%	8%	2%	11%
Smoked cigars, past 30 days	9%	3%	6%	12%	16%	6%	12%
Used snus, ever	8%	6%	6%	8%	12%	3%	12%

<sup>\*</sup> Suppressed due to small cell sizes (≤ 5 students)

Among students who smoked in the past 30 days:

Usual source of cigarettes, past 30 days

	All		Grade				K
	2011	9	10	11	12	Female	Male
Bought them in a store or from a vending machine	14%	*	*	*	31%	12%	16%
Gave someone else money to buy them for me	31%	33%	34%	33%	27%	31%	31%
Borrowed (or bummed) them from someone else; Person 18 years or older gave them to me; took them from a store or family member	47%	57%	53%	56%	35%	51%	44%
Some other way	8%	*	*	*	*	*	*

Percent who tried to quit, past 12 months

	AII	Grade				Sex	
	2011	9	10	11	12	Female	Male
Tried to quit	48%	52%	69%	39%	43%	50%	46%

<sup>\*</sup> Suppressed due to small cell sizes (≤ 5 students)

	All		Gra	ade		Sex	
	2011	9	10	11	12	Female	Male
Were in the same room with someone who was smoking, past 7 days	41%	35%	36%	44%	48%	40%	41%
Were in the same car with someone who was smoking, past 7 days	32%	31%	29%	34%	35%	31%	33%
Were asked about smoking by a health professional, past 12 months	42%	35%	39%	44%	51%	44%	41%
Discussed dangers of smoking with parents or guardian, past 12 months	31%	42%	31%	29%	23%	31%	32%
Think that out of 100 Vermont high school students, 56 or more smoke cigarettes	12%	12%	12%	12%	12%	14%	10%

	All		Gra	ade		Sex	<
	2011	9	10	11	12	Female	Male
Used marijuana, ever	37%	21%	30%	40%	56%	34%	39%
Used marijuana before age 13	8%	8%	9%	8%	7%	5%	11%
Used marijuana, past 30 days	22%	14%	20%	24%	32%	19%	25%
Used marijuana on 3 to 9 of the past 30 days	5%	3%	5%	6%	6%	5%	5%
Used marijuana on 10 or more of the past 30 days	10%	5%	8%	10%	16%	6%	13%
Used marijuana on school property, past 30 days	5%	3%	6%	3%	8%	3%	7%

	All		Gra	ade		Sex	<b>(</b>
	2011	9	10	11	12	Female	Male
Used cocaine, past 30 days	2%	2%	2%	2%	4%	*	4%
Used inhalants, ever	7%	12%	7%	5%	6%	7%	7%
Used heroin, ever	2%	*	*	2%	4%	*	3%
Used methamphetamines, ever	3%	3%	2%	2%	5%	2%	4%
Used hallucinogens, ever	9%	6%	8%	7%	14%	5%	12%
Used the prescription stimulant Ritalin or Adderall not prescribed to them, ever	5%	4%	4%	5%	6%	4%	6%
Used a prescription pain reliever such as OxyContin, Vicodin, or other prescription pain reliever not prescribed to them, ever	10%	8%	9%	11%	13%	8%	12%
Used a prescription pain reliever or a prescription stimulant not prescribed to them, ever	11%	9%	10%	13%	14%	9%	13%
Were offered, sold, or given an illegal drug on school property, past 12 months	14%	11%	14%	17%	16%	11%	18%

<sup>\*</sup> Suppressed due to small cell sizes (≤ 5 students)

## Attitudes and Perceptions about Alcohol, Cigarette, and Marijuana Use

Percent of students who think their parents would think it is wrong or very wrong for them to:

	All		Gra	Sex			
	2011	9	10	11	12	Female	Male
Smoke cigarettes	90%	97%	91%	89%	83%	91%	90%
Drink alcohol	72%	86%	74%	71%	58%	75%	70%
Smoke marijuana	82%	92%	82%	81%	75%	85%	79%

Percent of students who think it would be wrong or very wrong for someone their age to:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Smoke cigarettes	75%	85%	81%	71%	60%	78%	72%
Drink alcohol	49%	67%	55%	43%	32%	52%	47%
Smoke marijuana	60%	74%	63%	54%	48%	67%	52%

## Attitudes and Perceptions about Alcohol, Cigarette, and Marijuana Use

Percent of students who think people their age greatly risk harming themselves (physically or in other ways) if they:

	All		Gra	Sex			
	2011	9	10	11	12	Female	Male
Smoke one or more packs of cigarettes per day	58%	59%	59%	58%	58%	64%	53%
Have five or more drinks of alcohol once or twice each weekend	40%	47%	39%	38%	36%	48%	32%
Smoke marijuana regularly	34%	50%	35%	28%	25%	41%	28%

Percent of students who report that it would be easy or very easy to get:

	All		Gra	Sex			
	2011	9	10	11	12	Female	Male
Cigarettes	63%	46%	54%	69%	83%	60%	67%
Alcohol	71%	59%	70%	74%	81%	70%	73%
Marijuana	57%	38%	52%	63%	75%	53%	61%

## **Sexual Behavior and Orientation**

	All	Gr	ade	Sex		
	2011	9 and 10	11 and 12	Female	Male	
Had sexual intercourse, ever	39%	23%	54%	39%	38%	
First had sexual intercourse before age 13	4%	3%	4%	2%	5%	
Had sexual intercourse with four or more people during their life	10%	6%	15%	10%	10%	
Had sexual intercourse at least once in the past three months	30%	17%	42%	32%	28%	
Had oral sex, ever	41%	29%	54%	43%	40%	
Had a HIV test, ever	11%	8%	14%	14%	8%	

### **Sexual Behavior and Orientation**

Among students who had sex in the past 3 months:

Percent of students who:

	All	Gr	ade	Sex		
	2011	9 and 10 11 and 12		Female	Male	
Used drugs or alcohol before last sexual intercourse	17%	21%	16%	8%	27%	
Used a condom for last sexual intercourse	60%	68%	58%	59%	62%	

Method used to prevent pregnancy during last sexual intercourse

	All	Gr	ade	Sex		
	2011	9 and 10 11 and 12		Female	Male	
Condoms	38%	47%	34%	31%	44%	
Prescription birth control including pills, rings, implants, injections,						
IUDs	48%	37%	52%	59%	37%	
No method	6%	10%	4%	4%	8%	
Withdrawal	5%	*	5%	4%	5%	
Some other method/not sure	3%	*	4%	*	6%	

<sup>\*</sup> Suppressed due to small cell sizes (≤ 5 students)

### **Sexual Behavior and Orientation**

Sexual orientation

	Sex	
	Female	Male
Heterosexual	92%	95%
Gay or lesbian	*	*
Bisexual	5%	2%
Not sure	3%	2%

With whom have you had sexual contact?

	Sex				
	Female Male				
Females	*	50%			
Males	51%	*			
Both	5%	3%			
Never had sexual contact	42%	46%			

<sup>\*</sup> Suppressed due to small cell sizes (≤ 5 students)

## **Body Image**

Percent of students who are:

	All	Grade			Sex		
	2011	9	10	11	12	Female	Male
Overweight (85th BMI Percentile)	15%	15%	19%	15%	12%	15%	16%
Obese (95th BMI Percentile)	10%	6%	12%	12%	12%	6%	15%

BMI=body mass index; weight in kilograms divided by height in meters, squared

Percent of students who think that they are:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Underweight	12%	15%	9%	9%	13%	8%	15%
About the right weight	61%	66%	57%	59%	60%	61%	60%
Overweight	28%	18%	34%	31%	27%	31%	25%

# **Body Image**

Percent of students who are trying to:

	All	Grade			Sex		
	2011	9	10	11	12	Female	Male
Lose weight	41%	33%	44%	43%	42%	51%	31%
Gain weight	11%	12%	9%	13%	11%	3%	20%
Stay the same weight	25%	29%	24%	23%	23%	26%	23%
Not trying to do anything about weight	23%	26%	23%	20%	24%	20%	26%

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Skipped meals to lose weight or keep from gaining weight, past 30 days	18%	15%	20%	18%	20%	26%	11%
Used an indoor tanning device such as a sunlamp, sunbed, or tanning booth, past 30 days	9%	7%	6%	12%	14%	15%	4%

## **Nutrition**

Percent of students who, during the past seven days:

	All		Gra	ade		Sex	x
	2011	9	10	11	12	Female	Male
Ate fruits and vegetables five or more times per day	24%	25%	24%	23%	25%	22%	26%
Ate two or more servings of fruit or fruit juice per day	34%	34%	36%	30%	37%	33%	35%
Ate three or more servings of vegetables per day	17%	15%	16%	17%	19%	14%	19%
Ate breakfast seven of the past seven days	45%	49%	44%	46%	38%	39%	50%
Ate breakfast at least three of the past seven days	80%	82%	79%	83%	76%	79%	81%
Ate breakfast zero of the past seven days	7%	7%	6%	7%	8%	7%	7%
Drank a can, bottle, or glass of soda or pop at least once per day	16%	15%	17%	16%	16%	10%	22%
Drank a can, bottle, or glass of a sugar-sweetened beverage such as lemonade, sweetened tea or coffee rinks, sports drinks, energy drinks, Snapple, or Sunny Delight at least							
once per day	15%	13%	13%	16%	16%	12%	17%

## **Physical Activity**

Percent of students who were physically active on:

	All	Grade			Sex		
	2011	9	10	11	12	Female	Male
Seven of the past seven days	26%	28%	30%	24%	24%	18%	34%
Zero of the past seven days	10%	7%	8%	13%	13%	13%	8%

Percent of students who attend physical education classes:

	All	Grade				Sex	
	2011	9	10	11	12	Female	Male
Every day during an average school week	13%	25%	14%	8%	8%	13%	14%
At least one day during an average school week	37%	66%	35%	21%	28%	37%	37%

Percent of students who, on an average school day, watch TV, play video or computer games, or use a computer for something that is not school work:

	All	Grade			Sex		
	2011	9	10	11	12	Female	Male
Three or more hours	33%	31%	33%	32%	35%	28%	36%
Five or more hours	8%	11%	8%	8%	7%	6%	11%

#### **Youth Assets**

Grades earned in school

	All	Grade			Sex		
	2011	9	10	11	12	Female	Male
Mostly As	39%	44%	40%	32%	42%	47%	32%
Mostly Bs	40%	39%	35%	47%	38%	37%	42%
Mostly Cs	14%	11%	16%	17%	12%	10%	18%
Mostly Ds or Fs	3%	2%	5%	2%	4%	3%	3%
None/not sure	4%	4%	4%	3%	4%	3%	4%

Percent of students who talk with their parents about school:

	All	Grade			Sex		
	2011	9	10	11	12	Female	Male
About every day	49%	52%	52%	46%	46%	52%	47%
About once or twice a week	29%	28%	28%	31%	27%	27%	30%
About once or twice a month	9%	7%	8%	10%	11%	9%	9%
Less than once a month	6%	6%	6%	6%	7%	6%	7%
Never	7%	7%	6%	7%	8%	7%	7%

#### **Youth Assets**

Percent of students who:

	All	Grade			Sex		
	2011	9	10	11	12	Female	Male
Volunteer one or more hours in an average week	58%	55%	60%	58%	61%	59%	58%
Volunteer three or more hours in an average week	20%	18%	21%	21%	19%	17%	22%

Percent of students who agree or strongly agree:

	All	Grade			Sex		
	2011	9	10	11	12	Female	Male
Teachers really care about you and give you a lot of encouragement	57%	59%	57%	54%	61%	55%	59%
Students help decide what goes on in your school	46%	53%	50%	40%	41%	47%	44%
In your community, you feel like you matter to people	53%	55%	56%	51%	52%	49%	58%

# 2011 Vermont Middle School Youth Risk Behavior Survey

### **Results for Addison County**

#### Participation

Gı	Grade Sex				
6 and 7	8	Female Male		All	
353	326	343	336	679	

Fewer than 25 Addison County sixth grade students were eligible and participated. Therefore, grades 6 and 7 were combined for this report.

For the middle school survey, we assigned county of residence based on the supervisory union/school district or, in some cases, individual schools.

#### **Participating Schools:**

Middlebury Union Middle School Mount Abraham Union MS/HS Orwell Village School Vergennes Union High School

# Demographics

Age

	All
	2011
12 or younger	32%
13	50%
14 or older	18%

# Race and Ethnicity

	All
	2011
White non-Hispanic	84%
Racial or ethnic minority	16%

Percent of students who:

	All	Grade		Sex	
	2011	6 and 7	8	Female	Male
Ride bicycles	92%	94%	90%	91%	92%

Of students who ride bicycles, percent who wear helmets:

	All	Grade		Sex	
	2011	6 and 7	8	Female	Male
Always	15%	15%	15%	17%	13%
Most of the time	23%	24%	22%	25%	21%
Sometimes	16%	14%	19%	17%	16%
Rarely or never	46%	47%	45%	42%	50%

Percent of students who:

	All	Grade		Sex	
	2011	6 and 7	8	Female	Male
Rollerblade or skateboard	42%	47%	36%	37%	47%

Of students who rollerblade or skateboard, percent who wear helmets:

	All	Grade		Sex	
	2011	6 and 7	8	Female	Male
Always	22%	21%	23%	26%	19%
Most of the time	20%	21%	18%	23%	17%
Sometimes	13%	15%	10%	13%	13%
Rarely or never	45%	43%	48%	38%	51%

Percent of students who wear seatbelts when riding in a car:

	All	Grade		Sex	
	2011	6 and 7	8	Female	Male
Always	70%	68%	71%	74%	65%
Most of the time	19%	20%	17%	18%	19%
Sometimes	6%	7%	6%	5%	8%
Rarely or never	5%	5%	6%	3%	8%

Percent of students who ever rode in a car driven by someone who had been drinking alcohol:

	All	Grade		Sex	
	2011	6 and 7	8	Female	Male
Yes	26%	25%	28%	26%	27%
No	54%	53%	54%	53%	54%
Not sure	27%	30%	25%	29%	26%

	All	Gr	ade	Sex	<
	2011	6 and 7	8	Female	Male
Were ever in a physical fight	49%	49%	48%	30%	68%
Did not go to school because felt unsafe, past 30 days	7%	7%	7%	8%	5%
Were ever bullied on school property	47%	45%	49%	46%	48%
Were ever electronically bullied	25%	19%	31%	34%	16%
Were bullied, past 30 days	30%	28%	31%	31%	28%
Bullied someone else, past 30 days	17%	16%	19%	15%	19%
Ever seriously thought about killing themselves	18%	14%	22%	21%	15%
Ever made a suicide plan	9%	6%	13%	13%	6%
Ever attempted suicide	3%	3%	4%	4%	3%

\_\_\_\_

# Alcohol, Tobacco, and Other Drug Use

	All Grade		Sex		
	2011	6 and 7	8	Female	Male
Ever had a drink of alcohol other than a few sips	24%	18%	30%	17%	31%
Had first drink other than a few sips before age 11	9%	9%	9%	6%	13%
Drank alcohol, past 30 days	9%	6%	12%	6%	12%
Had five or more drinks of alcohol in a row, past 30 days	3%	*	5%	2%	5%

<sup>\*</sup> Suppressed due to small cell sizes (≤ 5 students)

## Alcohol, Tobacco, and Other Drug Use

	All	Grade		Sex	
	2011	6 and 7	8	Female	Male
Tried cigarette smoking, even one or two puffs, ever	13%	11%	15%	10%	16%
Smoked before age 11	3%	3%	3%	*	4%
Smoked in past 30 days	4%	2%	5%	3%	5%
Used chewing tobacco, past 30 days	4%	2%	5%	*	6%
Smoked cigars, past 30 days	2%	*	2%	*	2%
Used snus, ever	3%	2%	4%	2%	4%

<sup>\*</sup> Suppressed due to small cell sizes (≤ 5 students)

# Alcohol, Tobacco, and Other Drug Use

	All	Grade		Sex	
	2011	6 and 7	8	Female	Male
Were in the same room with someone who was smoking, past 7 days	29%	28%	31%	29%	30%
Were in the same car with someone who was smoking, past 7 days	23%	23%	24%	23%	24%
Were asked about smoking by a health professional, past 12 months	20%	18%	23%	23%	18%
Discussed dangers of smoking with parents or guardian, past 12 months	43%	46%	40%	45%	41%
Think that out of 100 Vermont high school students, 56 or more smoke cigarettes	10%	10%	10%	10%	11%

# Alcohol, Tobacco, and Other Drug Use

	All	Grade		Sex	
	2011	6 and 7	8	Female	Male
Used marijuana, ever	5%	3%	8%	3%	7%
Used marijuana before age 11	*	*	*	*	*
Used marijuana, past 30 days	3%	*	4%	2%	4%
Used inhalants, ever	7%	8%	6%	8%	5%
Misused prescription drugs, ever	3%	2%	3%	*	4%

<sup>\*</sup> Suppressed due to small cell sizes (≤ 5 students)

#### Attitudes and Perceptions about Alcohol, Cigarette, and Marijuana Use

Percent of students who think their parents would think it is wrong or very wrong for them to:

	All	Grade		Sex	
	2011	6 and 7	8	Female	Male
Smoke cigarettes	95%	97%	93%	96%	94%
Drink alcohol	89%	91%	86%	90%	88%
Smoke marijuana	95%	97%	93%	97%	94%

Percent of students who think it would be wrong or very wrong for someone their age to:

	All	Grade		Sex	
	2011	6 and 7	8	Female	Male
Smoke cigarettes	93%	96%	89%	95%	90%
Drink alcohol	87%	90%	83%	89%	84%
Smoke marijuana	92%	96%	88%	94%	90%

## Attitudes and Perceptions about Alcohol, Cigarette, and Marijuana Use

Percent of students who think people their age greatly risk harming themselves (physically or in other ways) if they:

	All	Gr	Grade		K
	2011	6 and 7	8	Female	Male
Smoke one or more packs of cigarettes per day	72%	68%	76%	78%	66%
Have five or more drinks of alcohol once or twice each weekend	56%	56%	56%	62%	50%
Smoke marijuana regularly	67%	67%	67%	72%	62%

Percent of students who report that it would be easy or very easy to get:

	All	Gr	Grade		<
	2011	6 and 7	8	Female	Male
Cigarettes	25%	18%	32%	21%	28%
Alcohol	40%	33%	47%	40%	39%
Marijuana	13%	7%	20%	11%	15%

# **Body Image**

Percent of students who think that they are:

	All	Grade		Sex	
	2011	6 and 7	8	Female	Male
Underweight	15%	17%	13%	14%	16%
About the right weight	58%	59%	56%	58%	57%
Overweight	27%	25%	31%	28%	27%

Percent of students who are trying to:

	All	Grade		Sex	
	2011	6 and 7	8	Female	Male
Lose weight	39%	37%	41%	45%	32%
Gain weight	11%	11%	10%	6%	16%
Stay the same weight	25%	27%	22%	22%	28%
Not trying to do anything about weight	26%	25%	27%	28%	24%

	All	Grade 6 and 7 8		Sex	
	2011			Female	Male
Skipped meals to lose weight or keep from gaining weight, past 30					
days	20%	14%	25%	25%	14%

### Nutrition

Percent of students who ate breakfast on:

	All	Grade 6 and 7 8		Sex	(
	2011			Female	Male
Seven of the past seven days	46%	49%	43%	42%	51%
At least three of the past seven days	78%	80%	76%	75%	81%
Zero of the past seven days	10%	11%	9%	12%	8%

# **Physical Activity**

Percent of students who were physically active on:

	All	Grade		Grade		Sex	K
	2011	6 and 7 8		Female	Male		
Seven of the past seven days	35%	38%	32%	25%	45%		
Zero of the past seven days	6%	4% 8%		6%	7%		

Percent of students who attend physical education classes:

	All	Grade 6 and 7 8		Sex	<b>(</b>
	2011			Female	Male
Every day during an average school week	37%	37%	37%	36%	38%
At least one day during an average school week	90%	88%	91%	91%	89%

### **Physical Activity**

Percent of students who, on an average school day, watch TV:

	All	Grade		Sex	<
	2011	6 and 7 8		Female	Male
Three or more hours	22%	23%	22%	21%	24%
Five or more hours	6%	4% 7%		5%	7%

Percent of students who, on an average school day, play video or computer games or use a computer for something that is not school work:

	All	Grade		Sex	K
	2011	6 and 7 8		Female	Male
Three or more hours	21%	20%	23%	19%	24%
Five or more hours	7%	6%	8%	5%	10%

#### **Youth Assets**

Percent of students who, in the past 7 days, went online for something that was not for school:

	All	Grade		Sex	
	2011	6 and 7	8	Female	Male
Zero or one day	27%	34%	18%	23%	30%
Two or three days	18%	19%	18%	19%	17%
Four or five days	14%	12%	16%	15%	13%
Six or seven days	41%	34%	48%	43%	39%

Percent of students who talk with their parents about school:

	All	Grade		Sex	K
	2011	6 and 7 8		Female	Male
About every day	45%	41%	49%	46%	43%
About once or twice a week	34%	39%	28%	34%	33%
About once or twice a month	9%	9%	8%	8%	9%
Less than once a month	6%	5%	7%	6%	7%
Never	7%	6%	8%	6%	7%

#### **Youth Assets**

Percent of students who agree or strongly agree:

	All Grade Sex		Grade		<b>K</b>
	2011	6 and 7	8	Female	Male
Teachers really care about you and give you a lot of encouragement	60%	59%	62%	63%	58%
Students help decide what goes on in your school	44%	46%	43%	44%	45%
In your community, you feel like you matter to people	56%	58%	53%	57%	54%

# **Vermont High School Youth Risk Behavior Survey**

### **Recalculated Results for Addison County**

Before 2011, students in grades 8-12 took the Vermont Youth Risk Behavior Survey. In 2011, we conducted a high school survey of students in grades 9-12, and a middle school survey of students in grades 6-8.

All results in the 2011 high school survey report are for grades 9-12 only. The following pages contain rates from previous surveys recalculated for grades 9-12, overall and by gender, for comparison to 2011. Rates by grade did not change.

This section of the report includes only the variables on the 2011 YRBS. # = question was not asked that year.

### Personal Safety - All

			Υe	ear		
	1999	2001	2003	2005	2007	2009
Always/almost always wear seat belt when riding in car	80%	81%	87%	86%	83%	84%
Rode with drinking driver, past 30 days	27%	27%	20%	22%	24%	21%
Drove after drinking, past 30 days	9%	10%	6%	7%	9%	7%
Rode with marijuana user, past 30 days	28%	30%	23%	25%	22%	23%
Drive after marijuana use, past 30 days	14%	14%	11%	10%	10%	11%
Carried a weapon on school property, past 30 days	12%	11%	8%	11%	13%	12%
Did not go to school because felt unsafe, past 30 days	4%	4%	3%	3%	3%	4%
Threatened or injured with weapon on school property, past 12 months	7%	7%	6%	6%	5%	5%
In physical fight, past 12 months	25%	25%	22%	24%	24%	24%
In physical fight at school, past 12 months	14%	13%	12%	13%	12%	10%

### **Personal Safety - Females**

			Υe	ear		
	1999	2001	2003	2005	2007	2009
Always/almost always wear seat belt when riding in car	86%	85%	90%	89%	88%	88%
Rode with drinking driver, past 30 days	24%	26%	20%	22%	23%	21%
Drove after drinking, past 30 days	7%	6%	6%	4%	7%	5%
Rode with marijuana user, past 30 days	26%	27%	23%	23%	22%	22%
Drive after marijuana use, past 30 days	10%	11%	9%	7%	7%	8%
Carried a weapon on school property, past 30 days	4%	5%	3%	4%	4%	4%
Did not go to school because felt unsafe, past 30 days	4%	4%	4%	3%	3%	3%
Threatened or injured with weapon on school property, past 12 months	5%	7%	4%	4%	3%	4%
In physical fight, past 12 months	17%	18%	15%	17%	18%	18%
In physical fight at school, past 12 months	8%	9%	7%	8%	7%	6%

### **Personal Safety - Males**

			Υe	ear		
	1999	2001	2003	2005	2007	2009
Always/almost always wear seat belt when riding in car	74%	76%	84%	83%	79%	80%
Rode with drinking driver, past 30 days	30%	26%	20%	23%	24%	22%
Drove after drinking, past 30 days	12%	13%	6%	10%	10%	9%
Rode with marijuana user, past 30 days	30%	33%	23%	28%	22%	24%
Drive after marijuana use, past 30 days	18%	18%	13%	13%	12%	14%
Carried a weapon on school property, past 30 days	20%	16%	13%	20%	22%	20%
Did not go to school because felt unsafe, past 30 days	4%	3%	2%	3%	4%	4%
Threatened or injured with weapon on school property, past 12 months	9%	7%	7%	8%	7%	6%
In physical fight, past 12 months	32%	33%	28%	32%	31%	30%
In physical fight at school, past 12 months	20%	18%	16%	19%	17%	14%

### Personal Safety - All

	Year						
	1999	2001	2003	2005	2007	2009	
Were bullied, past 30 days	#	#	#	15%	15%	14%	
Bullied someone, past 30 days	#	#	#	19%	19%	18%	
Electronically bullied, past 12 months	#	#	#	#	#	13%	
Hit by boy/girlfriend on purpose, past 12 months	7%	8%	7%	5%	6%	7%	
Purposefully hurt self without wanting to die, ever	#	#	#	#	15%	14%	
Planned suicide, past 12 months	16%	15%	10%	11%	8%	7%	
Attempted suicide, past 12 months	7%	8%	6%	6%	2%	4%	

### **Personal Safety - Females**

	Year						
	1999	2001	2003	2005	2007	2009	
Were bullied, past 30 days	#	#	#	16%	15%	16%	
Bullied someone, past 30 days	#	#	#	14%	15%	15%	
Electronically bullied, past 12 months	#	#	#	#	#	18%	
Hit by boy/girlfriend on purpose, past 12 months	7%	8%	7%	4%	5%	5%	
Purposefully hurt self without wanting to die, ever	#	#	#	#	20%	19%	
Planned suicide, past 12 months	19%	19%	14%	13%	9%	9%	
Attempted suicide, past 12 months	9%	10%	8%	9%	3%	5%	

### **Personal Safety - Males**

	Year						
	1999	2001	2003	2005	2007	2009	
Were bullied, past 30 days	#	#	#	14%	15%	13%	
Bullied someone, past 30 days	#	#	#	25%	23%	21%	
Electronically bullied, past 12 months	#	#	#	#	#	8%	
Hit by boy/girlfriend on purpose, past 12 months	7%	8%	7%	7%	8%	8%	
Purposefully hurt self without wanting to die, ever	#	#	#	#	9%	8%	
Planned suicide, past 12 months	12%	10%	6%	7%	6%	5%	
Attempted suicide, past 12 months	4%	5%	3%	3%	2%	3%	

Alcohol - All

			Υe	ear		
	1999	2001	2003	2005	2007	2009
Consumed alcohol, ever	75%	70%	65%	65%	63%	64%
Drank alcohol prior to age 13	27%	25%	21%	20%	19%	20%
Consumed alcohol, past 30 days	47%	43%	37%	37%	38%	36%
Binged on alcohol, past 30 days	30%	27%	21%	21%	24%	21%
3 or more days drinking, past 30 days	28%	25%	18%	18%	20%	18%
10 or more days drinking, past 30 days	6%	6%	5%	4%	5%	4%
3 or more days binging, past 30 days	13%	11%	9%	6%	9%	9%
Drank on school property, past 30 days	5%	6%	3%	4%	3%	3%

#### **Alcohol - Females**

			Υe	ear		
	1999	2001	2003	2005	2007	2009
Consumed alcohol, ever	74%	71%	65%	63%	64%	65%
Drank alcohol prior to age 13	22%	23%	17%	17%	16%	16%
Consumed alcohol, past 30 days	46%	44%	39%	36%	38%	37%
Binged on alcohol, past 30 days	27%	26%	21%	18%	23%	20%
3 or more days drinking, past 30 days	23%	24%	17%	15%	19%	15%
10 or more days drinking, past 30 days	4%	5%	4%	3%	4%	2%
3 or more days binging, past 30 days	9%	8%	6%	4%	8%	7%
Drank on school property, past 30 days	2%	5%	2%	3%	3%	2%

#### **Alcohol - Males**

			Υe	ear		
	1999	2001	2003	2005	2007	2009
Consumed alcohol, ever	77%	69%	65%	67%	63%	64%
Drank alcohol prior to age 13	33%	27%	24%	24%	22%	23%
Consumed alcohol, past 30 days	48%	42%	34%	37%	38%	36%
Binged on alcohol, past 30 days	34%	29%	23%	24%	25%	22%
3 or more days drinking, past 30 days	34%	25%	19%	21%	22%	21%
10 or more days drinking, past 30 days	8%	8%	7%	6%	6%	6%
3 or more days binging, past 30 days	17%	13%	11%	8%	11%	11%
Drank on school property, past 30 days	7%	7%	4%	5%	4%	4%

#### Alcohol - All

Of students who drank in the past 30 days

			Ye	ear		
	1999	2001	2003	2005	2007	2009
Usually consumed liquor such as vodka, rum, scotch, bourbon, or	#	#	#	#	42%	440/
whiskey	#	#	#	#	42%	41%
Usually consumed beer	#	#	#	#	33%	33%
Usually consumed alcopops	#	#	#	#	19%	19%
Usually consumed wine or wine coolers	#	#	#	#	5%	7%
Someone gave alcohol to me	#	#	#	#	#	51%
Gave someone money to buy alcohol for me	#	#	#	#	#	26%
Got or stole alcohol from home	#	#	#	#	#	15%
Bought in a store	#	#	#	#	#	5%
Stole alcohol from a store or restaurant	#	#	#	#	#	3%

#### **Alcohol - Females**

Of students who drank in the past 30 days

			Ye	ear		
	1999	2001	2003	2005	2007	2009
Usually consumed liquor such as vodka, rum, scotch, bourbon, or whiskey	#	#	#	#	47%	38%
Usually consumed beer	#	#	#	#	22%	24%
Usually consumed alcopops	#	#	#	#	26%	29%
Usually consumed wine or wine coolers	#	#	#	#	6%	9%
Someone gave alcohol to me	#	#	#	#	#	58%
Gave someone money to buy alcohol for me	#	#	#	#	#	26%
Got or stole alcohol from home	#	#	#	#	#	13%
Bought in a store	#	#	#	#	#	2%
Stole alcohol from a store or restaurant	#	#	#	#	#	1%

#### **Alcohol - Males**

Of students who drank in the past 30 days

			Ye	ear		
	1999	2001	2003	2005	2007	2009
Usually consumed liquor such as vodka, rum, scotch, bourbon, or	#	#	#	#	200/	4.40/
whiskey	#	#	#	#	38%	44%
Usually consumed beer	#	#	#	#	46%	44%
Usually consumed alcopops	#	#	#	#	12%	7%
Usually consumed wine or wine coolers	#	#	#	#	4%	6%
Someone gave alcohol to me	#	#	#	#	#	44%
Gave someone money to buy alcohol for me	#	#	#	#	#	25%
Got or stole alcohol from home	#	#	#	#	#	17%
Bought in a store	#	#	#	#	#	8%
Stole alcohol from a store or restaurant	#	#	#	#	#	6%

Tobacco - All

			Υe	ear		
	1999	2001	2003	2005	2007	2009
Smoked a cigarette, ever	55%	44%	32%	30%	29%	31%
Smoked cigarettes, past 30 days	31%	24%	17%	17%	15%	19%
Smoked daily, past 30 days	12%	16%	6%	6%	6%	6%
Smoked first cigarette prior to age 13	25%	18%	13%	12%	12%	11%
Smoked 11 or more cigarettes on days smoked, past 30 days	5%	4%	3%	2%	3%	3%
Smoked 20 or more cigarettes on days smoked, past 30 days	2%	2%	1%	1%	1%	1%
Used chewing tobacco, past 30 days	4%	5%	3%	6%	10%	8%
Smoked cigars, past 30 days	14%	12%	8%	#	#	14%

#### **Tobacco - Females**

	Year						
	1999	2001	2003	2005	2007	2009	
Smoked a cigarette, ever	56%	47%	35%	29%	29%	32%	
Smoked cigarettes, past 30 days	33%	28%	19%	16%	15%	20%	
Smoked daily, past 30 days	11%	19%	6%	5%	6%	7%	
Smoked first cigarette prior to age 13	22%	18%	13%	9%	11%	10%	
Smoked 11 or more cigarettes on days smoked, past 30 days	4%	3%	2%	1%	2%	3%	
Smoked 20 or more cigarettes on days smoked, past 30 days	1%	1%	0%	0%	1%	1%	
Used chewing tobacco, past 30 days	1%	3%	1%	1%	4%	3%	
Smoked cigars, past 30 days	5%	6%	5%	#	#	9%	

#### **Tobacco - Males**

			Υe	ear		
	1999	2001	2003	2005	2007	2009
Smoked a cigarette, ever	54%	41%	30%	32%	30%	30%
Smoked cigarettes, past 30 days	30%	21%	15%	17%	15%	17%
Smoked daily, past 30 days	14%	13%	6%	7%	6%	6%
Smoked first cigarette prior to age 13	27%	19%	14%	15%	13%	12%
Smoked 11 or more cigarettes on days smoked, past 30 days	7%	5%	4%	3%	3%	3%
Smoked 20 or more cigarettes on days smoked, past 30 days	3%	2%	2%	1%	1%	2%
Used chewing tobacco, past 30 days	8%	7%	5%	10%	15%	13%
Smoked cigars, past 30 days	24%	19%	11%	#	#	19%

# Other Drug Use - All

			Υe	ear		
	1999	2001	2003	2005	2007	2009
Smoked marijuana, ever	49%	47%	41%	38%	35%	39%
Tried marijuana prior to age 13	12%	12%	10%	9%	9%	9%
Used marijuana, past 30 days	31%	30%	25%	23%	21%	24%
Used marijuana 10 or more times, past 30 days	13%	14%	10%	10%	8%	10%
Used marijuana on school property, past 30 days	9%	10%	6%	7%	5%	6%
Used cocaine, past 30 days	4%	4%	4%	4%	4%	4%
Use inhalants, ever	15%	14%	10%	6%	10%	9%
Used heroin, ever	3%	4%	3%	2%	3%	2%
Used methamphetamines, ever	10%	10%	7%	5%	3%	3%
Used hallucinogens, ever	21%	19%	12%	12%	9%	11%
Were offered, sold, or given drugs on school property, past 12 months	31%	29%	23%	25%	19%	18%

## **Other Drug Use - Females**

			Υe	ear		
	1999	2001	2003	2005	2007	2009
Smoked marijuana, ever	46%	46%	38%	34%	34%	38%
Tried marijuana prior to age 13	8%	9%	7%	6%	7%	6%
Used marijuana, past 30 days	26%	27%	22%	18%	19%	23%
Used marijuana 10 or more times, past 30 days	8%	10%	7%	6%	7%	8%
Used marijuana on school property, past 30 days	6%	8%	5%	5%	4%	4%
Used cocaine, past 30 days	2%	4%	4%	4%	3%	1%
Use inhalants, ever	16%	16%	11%	5%	10%	7%
Used heroin, ever	2%	4%	2%	1%	2%	1%
Used methamphetamines, ever	9%	10%	8%	5%	3%	1%
Used hallucinogens, ever	16%	17%	11%	8%	6%	7%
Were offered, sold, or given drugs on school property, past 12 months	24%	26%	19%	22%	17%	14%

## Other Drug Use - Males

			Υe	ear		
	1999	2001	2003	2005	2007	2009
Smoked marijuana, ever	54%	48%	45%	41%	36%	40%
Tried marijuana prior to age 13	17%	16%	12%	12%	11%	12%
Used marijuana, past 30 days	36%	33%	27%	28%	22%	26%
Used marijuana 10 or more times, past 30 days	18%	18%	13%	14%	10%	14%
Used marijuana on school property, past 30 days	13%	13%	8%	9%	6%	9%
Used cocaine, past 30 days	6%	5%	4%	5%	5%	6%
Use inhalants, ever	15%	12%	10%	7%	9%	10%
Used heroin, ever	4%	4%	3%	3%	4%	4%
Used methamphetamines, ever	12%	9%	6%	6%	4%	5%
Used hallucinogens, ever	26%	21%	13%	15%	11%	14%
Were offered, sold, or given drugs on school property, past 12 months	39%	32%	27%	27%	22%	22%

## Attitudes and Perceptions about Alcohol, Cigarette, and Marijuana Use - All

			Υe	ear		
	1999	2001	2003	2005	2007	2009
Parents would think it is wrong/very wrong for you to smoke	82%	85%	91%	89%	91%	90%
Parents would think it is wrong/very wrong for you to drink	73%	75%	78%	73%	76%	76%
Parents would think it is wrong/very wrong for you to smoke marijuana	86%	85%	88%	88%	89%	86%
Wrong/very wrong for someone your age to smoke	53%	61%	74%	77%	74%	70%
Wrong/very wrong for someone your age to drink alcohol	40%	42%	51%	46%	46%	44%
Wrong/very wrong for someone your age to smoke marijuana	50%	54%	60%	63%	64%	60%
Easy/very easy to obtain cigarettes	77%	70%	69%	72%	67%	66%
Easy/very easy to obtain alcohol	71%	72%	69%	72%	69%	67%
Easy/very easy to obtain marijuana	66%	64%	60%	63%	58%	59%

## Attitudes and Perceptions about Alcohol, Cigarette, and Marijuana Use - Females

			Υe	ear		
	1999	2001	2003	2005	2007	2009
Parents would think it is wrong/very wrong for you to smoke	82%	87%	91%	90%	93%	92%
Parents would think it is wrong/very wrong for you to drink	74%	77%	77%	76%	79%	79%
Parents would think it is wrong/very wrong for you to smoke marijuana	88%	87%	89%	90%	89%	88%
Wrong/very wrong for someone your age to smoke	51%	59%	73%	76%	77%	73%
Wrong/very wrong for someone your age to drink alcohol	40%	42%	49%	48%	47%	46%
Wrong/very wrong for someone your age to smoke marijuana	52%	57%	62%	67%	65%	64%
Easy/very easy to obtain cigarettes	79%	72%	71%	71%	66%	65%
Easy/very easy to obtain alcohol	72%	73%	71%	71%	71%	66%
Easy/very easy to obtain marijuana	63%	63%	61%	61%	59%	57%

## Attitudes and Perceptions about Alcohol, Cigarette, and Marijuana Use - Males

			Υe	ear		
	1999	2001	2003	2005	2007	2009
Parents would think it is wrong/very wrong for you to smoke	82%	84%	92%	89%	90%	89%
Parents would think it is wrong/very wrong for you to drink	72%	73%	80%	71%	74%	72%
Parents would think it is wrong/very wrong for you to smoke marijuana	85%	82%	88%	86%	89%	84%
Wrong/very wrong for someone your age to smoke	55%	63%	76%	77%	70%	66%
Wrong/very wrong for someone your age to drink alcohol	40%	43%	52%	44%	45%	42%
Wrong/very wrong for someone your age to smoke marijuana	48%	51%	59%	60%	62%	55%
Easy/very easy to obtain cigarettes	75%	68%	68%	74%	67%	68%
Easy/very easy to obtain alcohol	70%	71%	68%	73%	67%	69%
Easy/very easy to obtain marijuana	69%	64%	59%	65%	57%	60%

#### **Sexual Behavior and Orientation - All**

			Υe	ear		
	1999	2001	2003	2005	2007	2009
Had sexual intercourse, ever	36%	35%	34%	34%	38%	39%
Had sex, past 3 months	27%	25%	24%	25%	29%	#
Had sex before age 13	6%	5%	4%	3%	4%	5%
4 or more sexual partners in lifetime	10%	10%	7%	8%	9%	12%
3 or more sexual partners in last 3 months	4%	4%	2%	2%	4%	#
Used alcohol or drugs prior to last sex (among sexually active)	29%	27%	24%	23%	23%	21%
Used condom at last sex (among sexually active)	65%	60%	69%	69%	73%	67%
Heterosexual	#	#	#	94%	93%	93%
Gay or lesbian, bisexual, not sure	#	#	#	6%	7%	7%

#### **Sexual Behavior and Orientation - Females**

			Υe	ear		
	1999	2001	2003	2005	2007	2009
Had sexual intercourse, ever	35%	35%	35%	32%	38%	37%
Had sex, past 3 months	28%	28%	27%	24%	30%	#
Had sex before age 13	4%	4%	3%	2%	3%	3%
4 or more sexual partners in lifetime	9%	10%	7%	7%	8%	9%
3 or more sexual partners in last 3 months	3%	3%	2%	1%	3%	#
Used alcohol or drugs prior to last sex (among sexually active)	22%	23%	23%	19%	20%	18%
Used condom at last sex (among sexually active)	56%	49%	68%	65%	67%	64%
Heterosexual	#	#	#	93%	90%	92%
Gay or lesbian, bisexual, not sure	#	#	#	7%	10%	8%

#### **Sexual Behavior and Orientation - Males**

			Υe	ear		
	1999	2001	2003	2005	2007	2009
Had sexual intercourse, ever	38%	35%	33%	37%	37%	39%
Had sex, past 3 months	25%	22%	22%	25%	28%	#
Had sex before age 13	9%	7%	5%	5%	5%	8%
4 or more sexual partners in lifetime	11%	10%	7%	9%	11%	14%
3 or more sexual partners in last 3 months	5%	5%	2%	3%	5%	#
Used alcohol or drugs prior to last sex (among sexually active)	35%	31%	25%	26%	26%	25%
Used condom at last sex (among sexually active)	74%	71%	70%	74%	80%	70%
Heterosexual	#	#	#	95%	96%	95%
Gay or lesbian, bisexual, not sure	#	#	#	5%	4%	5%

# **Body Image - All**

			Υe	ear		
	1999	2001	2003	2005	2007	2009
BMI between 85th-95th percentile	16%	15%	14%	17%	14%	13%
BMI >95th percentile	10%	11%	12%	11%	13%	14%
Underweight	13%	14%	14%	12%	13%	13%
Right weight	53%	54%	57%	56%	58%	59%
Overweight	34%	31%	29%	32%	30%	28%
Trying to lose weight	44%	41%	42%	43%	42%	40%
Trying to gain weight	10%	11%	13%	11%	11%	11%
Trying to stay at current weight	18%	18%	16%	18%	20%	22%
Doing nothing about weight	28%	30%	28%	28%	27%	27%

## **Body Image - Females**

	Year						
	1999	2001	2003	2005	2007	2009	
BMI between 85th-95th percentile	14%	12%	11%	14%	14%	11%	
BMI >95th percentile	7%	7%	9%	10%	10%	9%	
Underweight	11%	10%	8%	8%	9%	10%	
Right weight	49%	52%	55%	53%	57%	60%	
Overweight	41%	38%	36%	40%	34%	30%	
Trying to lose weight	62%	57%	60%	59%	55%	52%	
Trying to gain weight	2%	3%	3%	2%	3%	3%	
Trying to stay at current weight	16%	17%	15%	18%	19%	22%	
Doing nothing about weight	21%	24%	22%	21%	23%	23%	

## **Body Image - Males**

			Υe	ear		
	1999	2001	2003	2005	2007	2009
BMI between 85th-95th percentile	19%	18%	16%	20%	14%	14%
BMI >95th percentile	12%	15%	16%	13%	16%	19%
Underweight	16%	19%	19%	17%	16%	16%
Right weight	57%	56%	58%	59%	59%	58%
Overweight	26%	25%	23%	24%	25%	26%
Trying to lose weight	25%	24%	26%	27%	26%	28%
Trying to gain weight	19%	20%	23%	20%	20%	19%
Trying to stay at current weight	20%	20%	18%	19%	21%	22%
Doing nothing about weight	36%	36%	34%	34%	33%	31%

#### **Nutrition - All**

	Year					
	1999	2001	2003	2005	2007	2009
2 or more servings of fruit per day, past week	44%	40%	42%	41%	36%	36%
3 or more servings of vegetables per day, past week	12%	9%	14%	16%	15%	16%
5 or more servings of fruit/vegetables per day, past week	21%	16%	27%	27%	24%	24%
Breakfast on 7 or more days, past week	#	#	#	#	43%	46%
Breakfast on 3 or more days, past week	#	#	#	#	77%	80%
Breakfast on 0 days, past week	#	#	#	#	10%	8%
1 or more servings of soda per day, past week	#	#	#	#	20%	19%

#### **Nutrition - Females**

	Year					
	1999	2001	2003	2005	2007	2009
2 or more servings of fruit per day, past week	45%	38%	42%	41%	38%	34%
3 or more servings of vegetables per day, past week	12%	9%	15%	17%	16%	14%
5 or more servings of fruit/vegetables per day, past week	21%	15%	26%	26%	25%	22%
Breakfast on 3 or more days, past week	#	#	#	#	77%	78%
Breakfast on 7 or more days, past week	#	#	#	#	39%	44%
1 or more servings of soda per day, past week	#	#	#	#	12%	12%

#### **Nutrition - Males**

	Year					
	1999	2001	2003	2005	2007	2009
2 or more servings of fruit per day, past week	42%	43%	42%	42%	35%	37%
3 or more servings of vegetables per day, past week	12%	9%	14%	16%	15%	18%
5 or more servings of fruit/vegetables per day, past week	20%	17%	28%	28%	23%	27%
Breakfast on 3 or more days, past week	#	#	#	#	77%	82%
Breakfast on 7 or more days, past week	#	#	#	#	47%	48%
1 or more servings of soda per day, past week	#	#	#	#	28%	27%

## **Physical Activity - All**

	Year					
	1999	2001	2003	2005	2007	2009
60+ minutes of physical activity every day, past 7 days	#	#	#	#	27%	23%
60+ minutes of physical activity on zero days, past 7 days	#	#	#	#	11%	13%
P.E. at least once per week	44%	47%	45%	42%	42%	40%
P.E. five days per week	18%	27%	29%	20%	23%	13%
Watched tv/used computer for fun, 3 or more hours per day	#	28%	31%	34%	30%	36%
Watched tv/used computer for fun, 5 or more hours per day	#	7%	8%	9%	7%	10%

# **Physical Activity - Females**

	Year					
	1999	2001	2003	2005	2007	2009
60+ minutes of physical activity every day, past 7 days	#	#	#	#	19%	17%
60+ minutes of physical activity on zero days, past 7 days	#	#	#	#	13%	14%
P.E. at least once per week	42%	42%	42%	42%	39%	37%
P.E. five days per week	18%	23%	27%	19%	21%	11%
Watched tv/used computer for fun, 3 or more hours per day	#	22%	24%	30%	25%	29%
Watched tv/used computer for fun, 5 or more hours per day	#	4%	4%	6%	5%	5%

## **Physical Activity - Males**

	Year					
	1999	2001	2003	2005	2007	2009
60+ minutes of physical activity every day, past 7 days	#	#	#	#	35%	28%
60+ minutes of physical activity on zero days, past 7 days	#	#	#	#	8%	12%
P.E. at least once per week	45%	53%	48%	42%	45%	43%
P.E. five days per week	18%	31%	31%	21%	24%	15%
Watched tv/used computer for fun, 3 or more hours per day	#	33%	37%	38%	36%	42%
Watched tv/used computer for fun, 5 or more hours per day	#	9%	11%	12%	10%	14%

#### Youth Assets - All

	Year					
	1999	2001	2003	2005	2007	2009
Earned mostly As	#	32%	32%	34%	37%	36%
Earned mostly Bs	#	42%	41%	41%	38%	42%
Earned mostly Cs	#	18%	19%	16%	16%	14%
Earned mostly Ds or Fs	#	5%	5%	4%	4%	4%
Not sure/not graded	#	3%	4%	5%	4%	4%
Parents talk to you about school at least once a week	74%	72%	77%	77%	80%	79%
Volunteer, 1 or more hours per week	#	45%	44%	45%	44%	44%
Volunteer, 3 or more hours per week	#	15%	13%	15%	14%	15%
Agree/strongly agree that students help decide what goes on at school	#	43%	46%	45%	48%	48%
Agree/strongly agree that you matter to your community	#	39%	45%	47%	50%	51%

#### **Youth Assets - Females**

	Year					
	1999	2001	2003	2005	2007	2009
Earned mostly As	#	41%	38%	42%	47%	44%
Earned mostly Bs	#	40%	41%	40%	36%	39%
Earned mostly Cs	#	14%	14%	11%	10%	11%
Earned mostly Ds or Fs	#	2%	4%	3%	3%	3%
Not sure/not graded	#	3%	3%	4%	4%	3%
Parents talk to you about school at least once a week	74%	73%	78%	78%	83%	80%
Volunteer, 1 or more hours per week	#	51%	48%	49%	48%	47%
Volunteer, 3 or more hours per week	#	17%	15%	17%	15%	17%
Agree/strongly agree that students help decide what goes on at school	#	43%	42%	43%	50%	50%
Agree/strongly agree that you matter to your community	#	35%	38%	44%	48%	50%

#### **Youth Assets - Males**

	Year					
	1999	2001	2003	2005	2007	2009
Earned mostly As	#	22%	27%	26%	27%	29%
Earned mostly Bs	#	44%	40%	41%	41%	44%
Earned mostly Cs	#	23%	24%	21%	23%	18%
Earned mostly Ds or Fs	#	8%	5%	6%	5%	4%
Not sure/not graded	#	3%	4%	6%	4%	5%
Parents talk to you about school at least once a week	76%	71%	75%	75%	77%	78%
Volunteer, 1 or more hours per week	#	40%	40%	41%	40%	41%
Volunteer, 3 or more hours per week	#	12%	11%	12%	13%	13%
Agree/strongly agree that students help decide what goes on at school	#	44%	49%	48%	47%	47%
Agree/strongly agree that you matter to your community	#	42%	51%	49%	52%	51%