

Finding Support

While frequent waking for baby is normal and healthy, it can be very difficult for parents.

Lack of sleep can make it feel hard to follow safe sleep practices. If you are struggling, it's important to reach out for support.

- Talk to your Pediatrician about strategies specific to your child's needs.
- Ask family members, friends, or someone you trust to watch baby while you nap or catch up on sleep.
- PCAVT's Parent Helpline 1-800-CHILDREN (244-5373).

FOR MORE INFORMATION

on how to create a safe sleep environment for your baby, visit

UVMHealth.org/SafeSleep

American Academy of Pediatrics

HealthyChildren.org

U.S. Consumer Product Safety Commission

CPSC.gov

Safe Infant Sleep

SafeInfantSleep.org

Charlie's Kids

CharliesKids.org

Kids for Cribs

CribsforKids.org

The Period of Purple Crying

PurpleCrying.info

Created in partnership by:















Vermont Chapter

INCORPORATED IN VERMONT

American Academy of Pediatrics





Baby sleeps alone on their back on a flat surface

Baby sleeps in a crib, bassinet, portable crib, or play yard *

Remove unsafe sleep products such as pillows, blankets, stuffed animals, lounger, or pacifier cord/attachment Sometimes babies can't be soothed and that's okay. It does not harm a child's development. If you are stressed or overwhelmed, take care of yourself first and let baby stay in the safety of the crib. Crying is not harmful if you need space.

Share your room,
NOT your bed

Keep your baby cozy by using a one piece sleeper/suit or sleep sack

aby ig a Offer a

pacifier



Room Sharing

Baby should share your room, not your bed.

Room sharing means keeping your baby's crib, play yard, or bassinet in your bedroom, close to your bed for at least the first 6 months. This makes it easier to comfort or feed your baby, and then place them in their own sleep space when you're ready to go to sleep.

If there is any possibility that you might fall asleep while your baby is in your bed, make sure there are no pillows, sheets, blankets or any other items that could cover your baby's face, head and neck, or overheat them. As soon as you wake up, be sure to move your baby to their own bed.

Avoid falling asleep with your baby in other spots, too. The risk of sleep-related infant death is higher when infants sleep with someone on a couch, soft armchair, or cushion.

It's extra important not to bed share with your baby if you have been drinking alcohol or used cannabis, illicit drugs, or any medications that cause drowsiness or impact sleep.

If your baby falls asleep in a car seat, stroller, swing, infant carrier or sling, move them to a firm sleep surface on their back as soon as possible.

* Consumer Product Safety Commission Approved