CAS 117-82-8 Di(methoxyethyl) phthalate (DMEP)



 $C_{14}H_{18}O_{6}$



Summary of Health Effects

DMEP can cause reproductive and developmental harm in animals and may harm humans. The European Union has banned DMEP in cosmetics and children's toys.

How is DMEP used?

DMEP was detected as a plasticizer in a polyvinyl chloride (PVC) medical device and in personal care products collected from Shanghai markets.^{1,2} DMEP has been detected in PVC children's toys and articles³ including play and exercise balls, hoppers, and inflatable water products in Australia and in T-shirts, diapers, and house carpets in Germany.⁴

Toxicity: What are its health effects?

The European Union added DMEP to the Candidate List of Substances of Very High Concern (SVHC) due to reproductive toxicity and banned DMEP for use in cosmetics and accessible components of children's toys.^{5,6} The Globally Harmonized System for labelling classified DMEP as category 1B for reproductive toxicity and assigned the risk phrase "may damage fertility or the unborn child" and "suspected of damaging fertility."⁷

Animal studies reported reproductive and developmental harm in rodents fed DMEP or methoxyacetic acid (MAA), a metabolite of DMEP.^{4,8,9}

Exposure: How can a person come in contact with it?

A person can come in contact with DMEP by breathing in contaminated air or dust or by skin contact with contaminated house dust or consumer products.

DMEP has been detected in house dust in Germany.¹⁰ A Hong-Kong biomonitoring study detected DMEP in blood plasma of the male population.¹¹

References

1. Gimeno, P., Thomas, S., Bousquet, C., Maggio, A., Civade, C., Brenier, C., Bonnet, P. (2014). Identification and quantification of 14 phthalates and 5 non-phthalate plasticizers in PVC medical devices by GC-MS. *Journal of Chromatography B*, 949–950, 99–108.

- 2. Bao, J., Wang, M., Ning X., Zhou, Y., He Y., Yang J., Gao, X., Li, S., Ding, Z., Chen, B. (2015). Phthalate concentrations in personal care products and the cumulative exposure to female adults and infants in Shanghai. *Journal of Toxicology and Environmental Health, Part A*. 78, 325–341.
- 3. Al-Natsheh, M., Alawi, M., Fayyad, M., Tarawneh, I. (2015). Simultaneous GC–MS determination of eight phthalates in total and migrated portions of plasticized polymeric toys and childcare articles. *Journal of Chromatography B*, 985, 103–109.
- 4. U.S. Department of Health and Human Services. *Hazardous substances data bank* (HSDB, online database). National Toxicology Information Program, National Library of Medicine, Bethesda, MD. 1993.
- 5. European Commission. *Endocrine disruptor priority list*. Retrieved from <u>ec.europa.eu/environment/chemicals/endocrine/strategy/substances_en.htm</u>
- 6. Danish EPA. (2013). *Phthalate strategy*; Danish Environmental Protection Agency: Copenhagen, Denmark. Retrieved from <u>www2.mst.dk/Udgiv/publications/2013/06/978-87-93026-22-3.pdf</u>
- 7. European Commission. *Table of harmonized entries in Annex VI to CLP*. Retrieved from <u>echa.europa.eu/information-on-chemicals/annex-vi-to-clp</u>
- 8. Furr J., Lambright C., Wilson V.S., Foster P.M., Gray Jr, L.E. (2014). A short-term in vivo screen using fetal testosterone production, a key event in the phthalate adverse outcome pathway, to predict disruption of sexual differentiation. *Toxicological Sciences.*, 140(2), 403-424.
- 9. Rasjad, C., Yamashita, K., Datu, A. R., Yasuda, M. (1991). Pattern of limb malformations in mice induced by methoxyacetic acid, *Hiroshima Journal of Medical Sciences*, 40, 93–99.
- 10. Kersten, W. & Reich, T. (2003). Schwerfluchtige organische umweltchemikalien in Hamburger hausstäben [in German]. *Reinhaltung der luft* 63, 85–91.
- 11. Wan, H.T., Leung, P.Y., Zhao, Y.G., Wei, X., Wong, M.H., Wong, C.K. (2013). Blood plasma concentrations of endocrine disrupting chemicals in Hong Kong populations. *Journal of Hazardous Materials*, 261, 763–769.