

# Packing Vaccines for Transport during Emergencies

## Be ready BEFORE the emergency

Equipment failures, power outages, natural disasters—these and other emergency situations can compromise vaccine storage conditions and damage your vaccine supply. **It's critical to have an up-to-date emergency plan with steps you should take to protect your vaccine.** In any emergency event, activate your emergency plan immediately. Ideally, vaccine should be transported using a portable vaccine refrigerator or qualified pack-out. However, if these options are not available, you can follow the emergency packing procedures for refrigerated vaccines below:

## 1 Gather the Supplies




### Hard-sided coolers or Styrofoam™ vaccine shipping containers

- Coolers should be large enough for your location's typical supply of refrigerated vaccines.
- Can use original shipping boxes from manufacturers if available.
- Do NOT use soft-sided collapsible coolers.



### Conditioned frozen water bottles

- Use 16.9 oz. bottles for medium/large coolers or 8 oz. bottles for small coolers (enough for 2 layers inside cooler).
- Do NOT reuse coolant packs from original vaccine shipping container, as they increase risk of freezing vaccines.
- Freeze water bottles (can help regulate the temperature in your freezer).
- Before use, you must condition the frozen water bottles. Put them in a sink filled with several inches of cool or lukewarm water until you see a layer of water forming near the surface of bottle. The bottle is properly conditioned if ice block inside spins freely when rotated in your hand (this normally takes less than 5 minutes).



### Insulating material — You will need two of each layer

- **Insulating cushioning material** – Bubble wrap, packing foam, or Styrofoam™ for a layer above and below the vaccines, at least 1 in thick. Make sure it covers the cardboard completely. Do NOT use packing peanuts or other loose material that might shift during transport.
- **Corrugated cardboard** – Two pieces cut to fit interior dimensions of cooler(s) to be placed between insulating cushioning material and conditioned frozen water bottles.



- **Temperature monitoring device** – Digital data logger (DDL) with buffered probe. Accuracy of  $\pm 1^{\circ}\text{F}$  ( $\pm 0.5^{\circ}\text{C}$ ) with a current and valid certificate of calibration testing. Pre-chill buffered probe for at least 5 hours in refrigerator. Temperature monitoring device currently stored in refrigerator can be used, as long as there is a device to measure temperatures for any remaining vaccines.

## Why do you need cardboard, bubble wrap, and conditioned frozen water bottles?

Conditioned frozen water bottles and corrugated cardboard used along with one inch of Insulating cushioning material such as bubble wrap keeps refrigerated vaccines at the right temperature and prevents them from freezing. **Reusing vaccine coolant packs from original vaccine shipping containers can freeze and damage refrigerated vaccines.**



U.S. Department of  
Health and Human Services  
Centers for Disease  
Control and Prevention

Distributed by

Visit [www.cdc.gov/vaccines/SandH](http://www.cdc.gov/vaccines/SandH)  
for more information, or your state  
health department.



CS249275-I August 2015

# Packing Vaccines for Transport during Emergencies

## 2 Pack for Transport

### Conditioning frozen water bottles (this normally takes less than 5 minutes)


- Put frozen water bottles in sink filled with several inches of cool or lukewarm water or under running tap water until you see a layer of water forming near surface of bottle.
- The bottle is properly conditioned if ice block inside spins freely when rotated in your hand.
- If ice “sticks,” put bottle back in water for another minute.
- Dry each bottle.
- Line the bottom and top of cooler with a single layer of conditioned water bottles.
- Do NOT reuse coolant packs from original vaccine shipping container.




8. Temperature Monitoring Device Display (on lid)



7. Conditioned Water Bottles




6. Cardboard Sheet




5. Bubble wrap, packing foam, or Styrofoam™



4. Vaccines, Diluents, and Temperature Monitoring Device Probe



3. Bubble wrap, packing foam, or Styrofoam™



2. Cardboard Sheet



1. Conditioned Water Bottles

**Close lid** – Close the lid and attach DDL display and temperature log to the top of the lid.

**Conditioned frozen water bottles** – Fill the remaining space in the cooler with an additional layer of conditioned frozen water bottles.

**Insulating material** – Another sheet of cardboard may be needed to support top layer of water bottles.

**Insulating cushioning material** – Cover vaccines with another 1 in. layer of bubble wrap, packing foam, or Styrofoam™

**Vaccines** – Add remaining vaccines and diluents to cooler, covering DDL probe.

**Temperature monitoring device** – When cooler is halfway full, place DDL buffered probe in center of vaccines, but keep DDL display outside cooler until finished loading.

**Vaccines** – Stack boxes of vaccines and diluents on top of insulating material.

**Insulating cushioning material** – Place a layer of bubble wrap, packing foam, or Styrofoam™ on top (layer must be at least 1 in. thick and must cover cardboard completely).

**Insulating material** – Place 1 sheet of corrugated cardboard over water bottles to cover them completely.

**Conditioned frozen water bottles** – Line bottom of the cooler with a single layer of conditioned water bottles.

### NOTE:

This pack-out can maintain appropriate temperatures for up to 8 hours, but the container should not be opened or closed repeatedly.

## 3 Arrive at Destination

**Before opening cooler** – Record date, time, temperature, and your initials on vaccine temperature log.

**Storage** – Transfer boxes of vaccines quickly to storage refrigerator.

**Troubleshooting** – If there has been a temperature excursion, contact vaccine manufacturer(s) and/or your immunization program before using vaccines. Label vaccines “Do Not Use” and store at appropriate temperatures until a determination can be made.